


Wild Boars Carry Antimicrobial Resistant Enterobacteriaceae of Zoonotic Importance

Zoi Athanasakopoulou¹, Antonia Touloudi¹ and Charalambos Billinis^{1,2*}

¹Faculty of Veterinary Science, University of Thessaly, 43100 Karditsa, Greece

²Faculty of Public and One Health, University of Thessaly, 43100 Karditsa, Greece

ISSN: 2576-9162



Keywords: Wild boars; β -lactamases; ESBL; Carbapenemases; Antimicrobial resistance; One health

Mini Review

Antibiotics belonging to the β -lactam class are the most commonly used antimicrobials in human and veterinary medicine [1]. Bacteria initially susceptible to β -lactams can develop resistance by modifying the β -lactams' targets (mutation or alteration of penicillin binding proteins), reducing cell permeability (downregulation of porins required for β -lactam entry), overexpressing efflux systems or producing β -lactamases [2]. β -lactamases are enzymes capable of hydrolyzing the amide bond of the β -lactam ring, therefore rendering the antimicrobial ineffective [3]. The emergence of Extended Spectrum Beta Lactamase (ESBL) and Carbapenemase-Producing Enterobacteriaceae (CPE) is causing an unprecedented public health crisis, leaving few treatment options and affecting not only hospitalized patients, but also long-term care facilities and the community [4].

Genes encoding ESBLs and carbapenemases are typically located on transferable plasmids, which can be mobilized between the same or different bacterial species [5]. Additionally, clonal and horizontal gene transfer have contributed to the dissemination of β -lactamase-mediated Antimicrobial Resistance (AMR) from humans to animals or the environment and vice versa [6]. Thus, the capacity of such AMR determinants to transmit among various hosts raises concerns on a One Health perspective.

Wild boar (*Sus scrofa*) has been identified as an important reservoir of zoonoses and food-borne pathogens [7-9]. Interest has been drawn to the species due to its "invasive" behavior, the great increase of its population densities, its omnivorous diet and close proximity to anthropized environments (farms and pastures) that could favor health risks resulting from the transmission of infections across the wildlife-domestic animal-human interface [10]. In recent years, research has been conducted to identify carriage of genes conferring resistance to human last-resort antibiotics, including ESBL and carbapenem-resistance genes, from wild boars. Results indicate the dissemination of public health significant antimicrobial resistance determinants, such as those responsible for fluoroquinolone and colistin resistance [11,12]. For instance, quinolone resistant *Escherichia coli* (*E. coli*) isolates harboring plasmid-mediated genes of the qnr group (qnrS1/qnrS3) have been identified in Poland [12].

Regarding ESBLs, their dispersion among wild boars has been documented in several European and African countries, while data from other regions are not available. The overall rate of ESBL-Enterobacteriaceae fecal carriage among the sampled animals in the existing studies has been described to vary widely from 0.9% in Poland to 15.9% in Italy and from 28.1% in Tunisia up to 44.4% in Algeria [11-14]. Variants of the CTX-M family predominate in most of the studies, with CTX-M-1 being the heretofore most prevalent ESBL enzyme in *E. coli* isolates reported from European countries and in particular from Germany, Portugal, Spain and Czech Republic [15-19]. On the other hand, variant CTX-M-15 has been identified as the most prevalent in both *E. coli* and *Klebsiella pneumoniae* strains in Algeria [13]. Other ESBLs

***Corresponding author:** Charalambos Billinis, Faculty of Veterinary Science, Faculty of Public and One Health, Greece

Submission:  January 26, 2022

Published:  February 4, 2022

Volume 8 - Issue 5

How to cite this article: Zoi Athanasakopoulou, Charalambos Billinis, Antonia Touloudi. Wild Boars Carry Antimicrobial Resistant Enterobacteriaceae of Zoonotic Importance. *Appro Poultry Dairy & Vet Sci* 9(1). APDV.000702. 2022.
DOI: [10.31031/APDV.2022.09.000702](https://doi.org/10.31031/APDV.2022.09.000702)

Copyright@: Charalambos Billinis, This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

have also been reported, though to a lesser extent, and include CTX-M-14, SHV-12, SHV-28 and TEM-52b [14,19,20]. Concerning the transposable elements responsible for the respective ESBL genes' diffusion, few information is available and describes the presence of class 2 integrons among two *E. coli* strains [19]. Finally, different sequence types and incompatibility groups of plasmids have been associated with ESBLs in wild boars in the aforementioned studies.

The overuse and misuse of β -lactam molecules in both human and veterinary medicine as well as the ability of antibiotic residues and of resistant bacteria originating from human or animal excrements to accumulate in numerous environmental niches have been highlighted as the main causes of ESBL carriage by wild boars [11]. ESBLs were more frequently isolated from the urban and peri-urban wild boars in Spain than from the forest ones and increased human population density led to an increase of blaTEM prevalence in wild boars in Italy [14,20]. These facts indicate the great influence of humans in AMR diffusion. CPE colonization in wild boars has only been described by a sole report in Algeria, to date [21]. Plasmid located blaOXA-48 was identified in ST635 *E. coli* strains, whereas the same OXA variant was chromosomally encoded in ST13 *Klebsiella pneumoniae*. Transmission to wild boars was associated with their feeding habits in the Algerian warm and humid habitat that favors the preservation of contaminating bacteria of human and animal origin.

Overall, it is evident that β -lactamase resistance is not confined to human and livestock settings but has also arisen in wild fauna. Natural habitats become increasingly invaded by humans, forcing wildlife into greater direct and indirect contact with them and ultimately elevating the risk of AMR transmission among populations [22], especially as regards commensal, ubiquitous bacteria such as Enterobacteriaceae. Further research is required to fully unveil the role of wild boars as reservoirs and amplifiers of β -lactamase genes and their capacity to diffuse AMR back to humans. Special attention should be given during wild boar hunting and manipulation of wild boar meat, since the rising, complex challenge of AMR now requires a holistic and multidisciplinary approach.

Acknowledgement

This work has been co-funded by the European Union and the General Secretariat for Research and Innovation, Ministry of Development & Investments, under the project «Novel technologies for surveillance and characterization of Extended-spectrum β -lactamase and Carbapenemase producing Enterobacteriaceae, in humans and animals (CARBATECH)» T2DGE-0944, of the Bilateral S&T Cooperation Program Greece–Germany 2017. This support is gratefully acknowledged.

Conflicts of Interest

The authors declare no conflict of interest.

References

- (2019) Antimicrobial consumption in the Eu and EEA: annual epidemiological report. European Centre for Disease Prevention and Control.

- King DT, Sobhanifar S, Strynadka NCJ (2017) The mechanisms of resistance to β -lactam antibiotics. In: Berghuis A, Matlashewski G, et al. (Eds.), Handbook of Antimicrobial Resistance. Springer, New York, USA, pp. 177-201.
- Majiduddin FK, Materon IC, Palzkill TG (2002) Molecular analysis of beta-lactamase structure and function. Int J Med Microbiol 292(2): 127-137.
- Tzouveleki LS, Markogiannakis A, Psychogiou M, Tassios PT, Daikos GL (2012) Carbapenemases in *Klebsiella Pneumoniae* and Other enterobacteriaceae: An Evolving crisis of global dimensions. Clinical Microbiology Reviews 25(4): 682-707.
- Rossolini GM, D Andrea MM, Mugnaioli C (2008) The spread of CTX-M-type extended-spectrum beta-lactamases. Clin Microbiol Infect 14 (Suppl 1): 33-41.
- Dzidic S, Bedeković V (2003) Horizontal gene transfer-emerging multidrug resistance in hospital bacteria. Acta Pharmacol Sin 24(6): 519-526.
- Meng XJ, Lindsay DS, Sriranganathan N (2009) Wild boars as sources for infectious diseases in livestock and humans. Philosophical Transactions of the Royal Society B: Biological Sciences 364 (1530): 2697-2707.
- Ahomaa FM (2019) Wild boar: a reservoir of foodborne zoonoses. Foodborne Pathog Dis 16(3): 153-165.
- Touloudi A, Valiakos G, Athanasiou LV, Birtsas P, Giannakopoulos A, et al. (2015) A serosurvey for selected pathogens in Greek European wild boar. Vet Rec Open 2(2): e000077.
- Miller RS, Farnsworth ML, Malmberg JL (2013) Diseases at the livestock-wildlife interface: status, challenges, and opportunities in the United States. Preventive Veterinary Medicine 110(2): 119-132.
- Selmi R, Tayh G, Srairi S, Mamlouk A, Chehida FB, et al. (2022) Prevalence, risk factors and emergence of extended-spectrum β -lactamase producing-, carbapenem- and colistin-resistant Enterobacteriales isolated from wild boar (*sus scrofa*) in Tunisia. Microbial Pathogenesis 163: 105385.
- Wasył D, Zając M, Lalak A, Skarżyńska M, Samcik I, et al. (2018) Antimicrobial resistance in *Escherichia coli* isolated from wild animals in Poland. Microbial Drug Resistance 24(6): 807-815.
- Bachiri T, Bakour S, Ladjouzi R, Thongpan L, Rolain JM, et al. (2017) High rates of CTX-M-15-producing *Escherichia Coli* and *Klebsiella Pneumoniae* in wild boars and barbary macaques in Algeria. Journal of Global Antimicrobial Resistance 8: 35-40.
- Formenti N, Calò S, Parisio G, Guarneri F, Birbes L, et al. (2021) ESBL/AmpC-producing *Escherichia Coli* in wild boar: epidemiology and risk factors. Animals (Basel) 11(7): 1855.
- Poeta P, Radhouani H, Pinto L, Martinho A, Rego V, et al. (2009) Wild boars as reservoirs of extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* of different phylogenetic groups. Journal of Basic Microbiology 49(6): 584-588.
- Rodríguez CP, Alt K, Grobbel M, Hammerl JA, Irrgang A, et al. (2020) Wildlife as sentinels of antimicrobial resistance in Germany? Front Vet Sci 7: 627821.
- Holtmann AR, Meemken D, Müller A, Seinige D, Büttner K, et al. (2021) Wild boars carry extended-spectrum β -lactamase- and AMPC-producing *Escherichia Coli*. Microorganisms 9(2): 367.
- Cristóvão F, Alonso CA, Igrejas G, Sousa M, Silva V, et al. (2017) Clonal diversity of extended-spectrum beta-lactamase producing *Escherichia coli* isolates in fecal samples of wild animals. FEMS Microbiology Letters 364(5): fnx039.
- Literak I, Dolejska M, Radimersky T, Klimes J, Friedman M, et al. (2010) Antimicrobial-resistant faecal *Escherichia coli* in wild mammals in

- central Europe: Multiresistant *Escherichia coli* producing extended-spectrum beta-lactamases in wild boars. *Journal of Applied Microbiology* 108(5): 1702-1711.
20. Darwich L, Seminati C, Olvera JRL, Vidal A, Aguirre L, et al. (2021) Detection of beta-lactam-resistant *Escherichia Coli* and toxigenic *clostridioides* difficile strains in wild boars foraging in an anthropization gradient. *Animals* 11(6): 1585.
21. Bachiri T, Bakour S, Lalaoui R, Belkebla N, Allouache M, et al. (2018) Occurrence of Carbapenemase-producing Enterobacteriaceae isolates in the wildlife: first report of oxa-48 in wild boars in Algeria. *Microbial Drug Resistance* 24(3): 337-345.
22. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, et al. (2008) Global trends in emerging infectious diseases. *Nature* 451(7181): 990-993.

For possible submissions Click below:

[Submit Article](#)